# V1309 Scorpii: merger of a contact binary \*

R. Tylenda<sup>1</sup>, M. Hajduk<sup>1</sup>, T. Kamiński<sup>1</sup>, A. Udalski<sup>2,3</sup>, I. Soszyński<sup>2,3</sup>, M. K. Szymański<sup>2,3</sup>, M. Kubiak<sup>2,3</sup>,
 G. Pietrzyński<sup>2,3,4</sup>, R. Poleski<sup>2,3</sup>, Ł. Wyrzykowski<sup>3,5</sup>, and K. Ulaczyk<sup>2,3</sup>

<sup>1</sup> Department for Astrophysics, N. Copernicus Astronomical Center, Rabiańska 8, 87-100 Toruń, Poland e-mail: tylenda, cinek, tomkam@ncac.torun.pl

<sup>2</sup> Warsaw University Observatory, Al. Ujazdowskie 4, 00-478 Warsaw, Poland

e-mail: udalski, soszynsk, msz, mk, pietrzyn, rpoleski, kulaczyk@astrouw.edu.pl

<sup>3</sup> The Optical Gravitational Lensing Experiment

<sup>4</sup> Universidad de Concepción, Departamento de Astronomia, Casilla 160–C, Concepción, Chile

<sup>5</sup> Institute of Astronomy, University of Cambridge, Madingley Road, Cambridge CB3 0HA, UK e-mail: wyrzykow@ast.cam.ac.uk

Received; accepted

#### ABSTRACT

*Context.* Stellar mergers are expected to take place in numerous circumstences in the evolution of stellar systems. In particular, they are considered as a plausible origin of stellar eruptions of the V838 Mon type. V1309 Sco is the most recent eruption of this type in our Galaxy. The object was discovered in September 2008.

Aims. Our aim is to investigate the nature of V1309 Sco.

*Methods.* V1309 Sco has been photometrically observed in course of the OGLE project since August 2001. We analyse these observations in different ways. In particular, periodogram analyses were done to investigate the nature of the observed short term variability of the progenitor.

*Results.* We find out that the progenitor of V1309 Sco was a contact binary with an orbital period of  $\sim$ 1.4 day. This period was decreasing with time. Similarly the light curve of the binary was also evolving, indicating that the system evolved toward its merger. The violent phase of the merger, marked by the systematic brightenning of the object, started in March 2008, i.e. half a year before the outburst discovery. We also investigate the observations of V1309 Sco during the outburst and the decline and show that they can be fully accounted for within the merger hypothesis.

*Conclusions.* For the first time in the literature we show, from direct observations, that contact binaries indeed end up by merging into a single object, as it was suggested in numerous theoretical studies of these systems. Our study also shows that stellar mergers indeed result in eruptions of the V838 Mon type.

Key words. stars: individual: V1309 Sco - stars: binaries: general - stars: peculiar

#### 1. Introduction

Stellar mergers have for a long time been recognized to play an important role in evolution of stellar systems. High stellar densities in globular clusters can lead to often collisions and mergers of stars (Leonard 1989). In this way the origin of blue strugglers can be explained. In dense cores of young clusters multiple mergers of protostars have been suggested as a way of formation of the most massive stars (Bonnell et al. 1998). Some binary stars, in particular contact binaries, are suggested to end up their evolution as stellar mergers (Robertson & Eggleton 1977).

The powerful outburst of V838 Mon in 2002 (Munari et al. 2002), accompanied by a spectacular light echo (Bond et al. 2003), raised interest in a class of stellar eruptions named "red novae", "optical transients" or "V838 Mon type eruptions". These objects, reaching typically a maximum luminosity of ~  $10^6 L_{\odot}$ , evolve to low effective temperatures and decline as very cool (super)giants. Apart form V838 Mon, in our Galaxy the class also includes V4332 Sgr, whose outburst was observed in 1994 (Martini et al. 1999), and V1309 Sco,

which erupted in 2008 (Mason et al. 2010). As extragalactic eruptions of this kind one can mention M31 RV (eruption in 1989, Mould et al. 1990), M85 OT2006 (Kulkarni et al. 2007), and NGC300 OT2008 (Berger et al. 2009).

Several interpretations of the eruptions were proposed. They include an unusual classical nova (Iben & Tutukov 1992; Shara et al. 2010), a late He-shell flash (Lawlor 2005), or a thermonuclear shell flash in an evolved massive star (Munari et al. 2005). Tylenda & Soker (2006) presented numerous arguments against these mechanisms. Instead, they showed that all the main observational characteristics of the V838 Mon type eruptions can be consistently understood as resulting from stellar collisions and mergers, as originally proposed in Soker & Tylenda (2003). For these reasons Soker & Tylenda (2007) proposed to term these kind of eruptions as mergebursts. Recently, Kashi et al. (2010) claimed that the V838 Mon type eruptions are of the same nature as some eruptions of luminous blue variables and that they can be powered by mass transfer events in binary systems. They however do not show, how a binary companion could transfer a mass of a few tens of solar mass onto a main sequence star primary on a time scale of  $\sim 2$  months, as it would be required to explain the eruption of V838 Mon (Tylenda & Soker 2006).

Send offprint requests to: R. Tylenda

<sup>\*</sup> Based on observations obtained with the 1.3-m Warsaw telescope at the Las Campanas Observatory of the Carnegie Institution of Washington.

In the present paper, we show that the recent red nova, V1309 Sco, is a rosetta stone in the studies of the nature of the V838 Mon type eruptions. Archive photometric data collected for the object in the OGLE project during  $\sim$ 6 years before the outburst allow us to conclude that the progenitor of V1309 Sco was a contact binary. The system quickly evolved toward its merger, which resulted in the eruption observed in 2008.

# 2. V1309 Scorpii


V1309 Sco, also known as Nova Sco 2008, was discovered on 2.5 September 2008 (JD 2454712) (Nakano 2008). The subsequent evolution however showed that, as pointed out in Mason et al. (2010), this was not a typical classical nova. When declining the object evolved to very low effective temperatures and 8 months after discovery it was observed as a late M-type giant. Thus the object shares the principal characteristics of the V838 Mon type eruptions (Tylenda & Soker 2006). Other common features of V1309 Sco and the V838 Mon type eruptions include: outburst time scale of order of months, outburst amplitude (maximum minus progenitor) of 7 – 10 magnitudes, expansion velocities of a few hunderd km s<sup>-1</sup> (rather than a few thousands as in classical novae), and oxide bands observed in later epochs implying oxygen rich (C/O < 1) matter involved in the eruption.

## 3. Observations

Owing to the position of V1309 Sco close to the Galactic centre  $(1 = 359^{\circ}8, b = -3^{\circ}1)$  the object appears to be situated within a field monitored in the OGLE-III and OGLE-IV projects (Udalski 2003)<sup>1</sup>. As a result V1309 Sco was observed on numerous occasions since August 2001. Altogether more than 2000 measurements, predominantly in the *I* Cousins photometric band, were obtained. Among them, ~1340 observations were done prior to the discovery of the object as a nova in September 2008. A few observations were also done in the *V* band (7 prior the discovery). The data were reduced and calibrated using standard OGLE procedures (Udalski et al. 2008). A typical precision of the measurements was 0.01 magnitude.

# 4. Results

All the measurements of V1309 Sco derived from the OGLE-III and IV surveys in the I photometric band are displayed in Fig. 1. The gaps in the data are due to conjunctions of the object with the Sun. Apart from 2001 and 2009 most of the data were obtained between February and October of each year. In 2001 the OGLE-III project was just starting to operate, hence only 11 measurements were obtained in this year (they were omitted from the analysis discribed below). In the period 2002–2008, from 52 (2002, 2003) to 367 (2006) observations were done each year. Near the maximum of the 2008 eruption, i.e. when the object was brighter than  $I \simeq 11$ , its image was overexposed in the OGLE frames, hence no data for this period (a few open points in Fig. 1 in this period display AAVSO results<sup>2</sup>). In May 2009 OGLE-III phase ended and threfore only 64 data points were collected during the 2009 observing season. OGLE project resumed regular observations of the Galactic center in March 2010 with a new instrumental setup, namely a 32 chip mosaic camera, starting OGLE-IV phase. 655 measurements were obtained in 2010.



**Fig. 2.** The evolution of the period of the photometric variations of the V1309 Sco progenitor. The line shows a least square fit of an exponential formula to the data (see text and Eq. 1).

As can be seen from Fig. 1, the progenitor of V1309 Sco was initially slowly increasing in brightness on a time scale of years and reached a local maximum in April 2007. Subsequently, during a year, the star faded by ~1 magnitude. In March 2008, the main eruption started, which led to the object discovery, as Nova Sco 2008, half a year later. Remarkable is the smooth, roughly exponential rise in brightness. During this event the object brightened by ~10 magnitudes, i.e. by a factor of  $10^4$ . An analysis of the data for the progenitor (2002–2007) is presented in Sect. 5, while the outburst and the decline of the object are discussed in Sect. 6.

# 5. The progenitor


# 5.1. The data

A short term variability, resulting in a ~0.5 magnitude scatter of the points in Fig. 1, is the most remarkable and interesting feature of the V1309 Sco progenitor. This variability is stricktly periodic. To show this, we have used a method employing periodic orthogonal polynomials to fit the observations and an analysis of variance to evaluate the quality of the fit, as described in Schwarzenberg-Czerny (1996). This method is particularly suitable for analysing unevenly sampled data of non-sinusoidal periodic variations, such as eclipses in binary systems. The resultant periodograms obtained for the particular observational epochs are presented in Appendix(A). They show that the principal periodicity in the variations observed in epochs 2 - 7 corresponds to a period of ~1.4 day. However, the derived period was not constant, but slowly decreasing with time, as presented in Fig. 2. It decreased by 1.2% during the pre-outburst observations.

When deriving the period plotted in Fig. 2 we devided the data sequence available for a given epoch into subsamples, each containing  $\sim$ 50 data points, and constructed periodograms for each subsample separately. In this way we can see that during

<sup>&</sup>lt;sup>1</sup> http://ogle.astrouw.edu.pl

<sup>&</sup>lt;sup>2</sup> available at http://www.aavso.org/



**Fig. 1.** The light curve of V1309 Sco: *I* magnitude versus time of observations in Julian Dates. Full points: data from the OGLE-III and OGLE-IV projects. Open points: data obtained by the AAVSO team during the 2008 eruption. The epochs, analysed in the text and in Fig. 3, are numbered with the last digit of the year of observations (e.g. 4 marks the data obtained in 2004).

later epochs, particularly in epochs 6 and 7, the period was significantly varying on a time scale of months.


We find that the time evolution of the observed period can be fitted by an exponential formula. A result of a least square fit (inverse errors used as weights) is shown with the full line in Fig. 2. The obtained formula is

$$P = 1.4456 \exp\left(\frac{15.29}{t - t_0}\right),\tag{1}$$

where *P* is the period in days, *t* is the time of observations in Julian Dates, and  $t_0 = 2455233.5$ .

Using the above period fit we folded up the observations and obtained light curves of the object in particular epochs. The results are displayed in Fig. 3. In 2002 - 2006 the light curves were obtained from all the data available for a particular epoch. The results are shown in the upper part of Fig. 3. In 2007 the light curve significantly evolved during the epoch and therefore, in the lower part of Fig. 3, we plotted the results for the subsamples (the same as when deriving the period) of epoch 7 (time goes from subsample a to e in the figure).

As can be seen from Fig. 3, the light curve in 2002 - 2006 displays two maxima and two minima. They are practically



**Fig. 3.** The light curves obtained from folding the data with the period described by Eq. (1). Upper part: epochs 2 - 6. Lower part: epoch 7 devided into five subsamples (time goes from a to e). The zero point of the magnitude (ordinate) scale is arbitrary.

equally spaced in the phase and, in the early epochs, have similar shapes. This is the reason why the periodograms for these epochs (see Appendix A) show a strong peak at a frequency twice greater (period twice shorter) than the main peak. In other words, the observations could have been interpreted with a period of ~0.7 day and a light curve with one maximum and minimum. The periodograms however show, already from epoch 2, that the period twice longer (~1.4 day) better reproduces the observations. The reason is seen in the later epochs, when the first maximum (at phase 0.25 in Fig. 3) in the light curve becomes increasingly stronger than the second one. As a result the peak

corresponding to the 0.7 day period deacreases and in epoch 7 it practically disappears.

#### 5.2. Interpretation

In principle one can consider three possible interpretations of the observed light variability, i.e. stellar pulsation, single star rotation, and eclipsing binary system.

### 5.2.1. Pulsation

The light curve in early epochs may imply a stellar pulsation with a period of  $\sim 0.7$  day and an amplitude of  $\sim 0.15$  magnitude. This interpretation, however, encounters severe problems when explaining the observed evolution of V1309 Sco. As we show below, the progenitor was probably a K-type star. There is no class of pulsating stars, which could be reconciled with these characteristics, i.e a K-type star pulsating with the above period and amplitude. Moreover, to explain the observed evolution of the light curve displayed in Fig. 3, one would have to postulate a switch of the star pulsation to a period exactly twice longer on a time scale of a few years, together with a gradual shortening of the period. This would be very difficult, if not impossible, to be understood within the present theory of stellar pulsation. Finally, there is no physical mechanism involving or resulting from a pulsation instability, which could explain the powerful 2008 eruption, when the object brightened by a factor of  $10^4$ .

#### 5.2.2. Rotation of a single star

The observed light curve could have been explained by a single K-type star rotating with a period of  $\sim 1.4$  day and having two spots (or rather two groups of spots) on its surface in early epochs. The two spots would be replaced by one spot in epoch 7. The object would thus belong to the FK Com class of rapidly rotating giants (e.g. Bopp & Stencel 1981). There are, however, several differences between the progenitor of V1309 Sco and the FK Com stars. The 1.4 day period is short as for the FK Com stars. The very good phase stability of the observed light curve. seen in Fig. 3, implies that the spot(s) would have had to keep the same position(s) on the star surface over a time span of a few years. This seems to be very improbable given differential rotation and meridional circulation expected to be substantial in the case of a rapidly rotating star. Indeed, in the case of the FK Com stars, the spots are observed to migrate and change their position on much shorter time scales (Korhonen et al. 2007). Also the systematic decrease of the rotational period is difficult to explain in the case of a single star, and is not observed in the FK Com stars. Finally, no eruption, as that of V1309 Sco in 2008. was observed for the FK Com stars. In fact, there is no known mechanism, which could produce such a huge eruption in the case of a single giant, even if it is fast rotating.

#### 5.2.3. A contact binary evolving to its merger

We are thus left with an eclipsing binary system. As we show below, this possibility allows us to explain all the principal characteristics of the observed evolution of the V1309 Sco progenitor, as well as the 2008 outburst.

The shape of the light curve, especially in early epochs (two rounded maxima of comparable brightness and two equally spaced minima), implies that the progenitor of V1309 Sco was a contact binary. The orbital period of  $\sim$ 1.4 day does not allow

-0.3

-0.2

-0.1

us to classify the object as a W UMa type binary, as the clasical W UMa stars have periods < 1 day. Nevertheless contact binaries with periods beyond 1 day are also observed (Paczyński et al. 2006). The exponentially decreasing period (Fig. 2) can be interpreted as resulting from an unstable phase of evolution of the system leading to the shrinkage of the binary orbit and finally to the merger of the system. Such a situation was predicted in theoretical studies (Robertson & Eggleton 1977; Rasio 1995) and it is now commonly believed that the W UMa binaries end their evolution by merging into a single star. Dissipation of the orbital energy in the initial, violent phase of the merger (Soker & Tylenda 2006) resulted in the V1309 Sco eruption observed in 2008.

One of the possible ways, which can lead to a merger of a binary system is the so-called Darwin instability. This happens, when the spin angular momentum of the system is more than a third of the orbital angular momentum. As a result, tidal interactions in the system cannot maintain any more the primary component in synchronization. The orbital angular velocity is higher than the primary's angular velocity, tidal forces increase their action and rapidly transport angular momentum from the orbital motion to the primary's rotation. In the case of contact binaries this takes place, when the binary mass ratio  $q \equiv M_2/M_1 \leq 0.1$  (Rasio 1995).

A maximum in the light curve of a typical contact binary is observed, when we look at the system more or less perpendicularly to the line joining the two components. As a result, we observe two maxima during each orbital revolution. The maxima are of similar brightness, as the system looks similar from both sides. The situation changes, when the Darwin instability starts working. The secondary starts orbiting faster than rotates the primary's envelope. The stars are in contact, so the difference in velocities is partly dissipated near the contact between the components. This should lead to formation of a brighter (hotter) region seen when we look at the leading side of the secondary. The system is then brighter than when we look from the opposite direction. The maxima in the light curve start differing. This is what we observe in the case of the V1309 Sco progenitor. As can be seen from Fig. 3, the first maximum (at phase  $\sim 0.25$ ) becomes progressively stronger than the second one (phase  $\sim 0.75$ ), when going from epoch 2 to epoch 6. In epoch 7, the second maximum disappears and, at the end of this epoch, we observe a light curve with only one maximum and one minimum. Apparently the system evolved to a more spherical configuration with a bright (hot) spot covering a large fraction of the system surface along the orbital plane.

#### 5.3. Basic parameters of the progenitor

# 5.3.1. Interstellar reddening, spectral type, and effective temperature

The effective temperature of the progenitor can be derived from the V - I colour. In epoch 6 a few measurements were obtained in the V band. They are displayed in Fig. 4 together with the I data obtained in the same time period (JD 2453880 – 2453910).

In order to determine the effective temeprature from a colour, an estimate of the interstellar reddening is also necessary. Mason et al. (2010) obtained  $E_{B-V} \simeq 0.55$  from interstellar lines observed in their spectra of V1309 Sco in outburst. We constrained spectral types of the object in outburst from the UVES spectra of Mason et al. (2010) obtained in their epochs 1 – 3 (downloaded from the VLT archive). Comparing them with the *BVRI* photometry of the object obtained by the

0.1 0.1 0.2 0.3 0.25 0.5 0.75 1 1.25 1.5 1.75 2 phase

**Fig. 4.** The V (open points) and I (asterisks) measurments obtained in JD 2453880 – 2453910 (epoch 6) and folded with the period described by Eq. (1). The zero point of the magnitude (ordinate) scale is arbitrary.

AAVSO team near the above spectral observations, we obtained 0.7  $\leq E_{B-V} \leq$  0.9. The OGLE data also allowed us do derive a *V* – *I* colour, equal to 5.27, 223 days after the outburst discovery date. This happened to be in the epoch of the SOAR spectroscopy of Mason et al. (2010), from which these authors derived a spectral type of M6–7. To reconcile the colour with the spectral type,  $E_{B-V} > 0.7$  is required. In the following we assume that V1309 Sco is reddened with  $E_{B-V} \approx 0.8$ .


As can be seen from Fig. 4, two *V* measurements were done close to the first maximum in the light curve (phase ~0.26), two others near the following minimum (phase ~0.55). The resultant colour, V - I, was 2.09 and 2.21 for the two phases, respectively. The mean value of the observed colour, i.e.  $V - I \approx 2.15$ , corrected for the reddening and compared with the standard colours of giants (luminosity class III) results in a spectral class K1–2 and an effective temperature  $T_{\rm eff} \approx 4500$  K for the progenitor of V1309 Sco.

The difference in the colour, as mentiond above, implies that at the maximum the object was on average  $\sim 200$  K hotter than in the minimum. This agrees with the interpretation of the evolution of the light curve done in Sect. 5.2.3.

#### 5.3.2. Luminosity and distance

The hypothesis that the progenitor of V1309 Sco was a contact binary allows us to evaluate other parameters of the system. Adopting a total mass of the binary,  $(M_1 + M_2)$ , of  $1.0 - 3.0 \text{ M}_{\odot}$  (observed range of masses of W UMa binaries) and an orbital period of 1.43 day, one obtains, from the third Kepler's law, that the separation of the components, *A*, is  $(3.7 - 5.4) \times 10^{11}$  cm (5.4  $- 7.7 \text{ R}_{\odot}$ ).

Taking the mass ratio, q, in the range 0.07 - 1.0 (at  $q \approx 0.07$  the Darwin instability is expected to set up in the W UMa systems, Jiang et al. 2010) we can derive the radii of the Roche



lobes of the components (Eggleton 1983). Assuming that both components fill their Roche lobes a total effective surface of the system can be obtained. This with the above estimate of  $T_{\rm eff}$  results in a luminosity of the system to be within  $(1.15 - 3.3) \times 10^{34}$  erg s<sup>-1</sup>  $(3.0 - 8.6 L_{\odot})$ .

Taking into account bolometric correction, interstellar reddening, and the mean observed brightness of the V1309 Sco progenitor in epoch 6, V = 18.22, we derive a distance to the object of  $3.0 \pm 0.7$  kpc. If the outer Roche lobes (dimensions can be calculated from Yakut & Eggleton 2005) are assumed to be filled by the components, the luminosity increases by ~35%, while the distance becomes  $3.5 \pm 0.7$  kpc. Note that the distance estimate is almost independent of the reddening value.

The above estimates of the luminosity and effective temperature, when compared to theoretical tracks of stellar evolution (e.g. Girardi et al. 2000), are consistent with a  $\sim 1.0 \text{ M}_{\odot}$  star at the beginning of the red giant branch.

#### 6. The outburst

#### 6.1. The data


Figure 5 (upper part) displays the rise of V1309 Sco during its eruption in 2008. This phase started in March 2008 and it took the object ~6 months to reach the maximum brightness. The rise was remarkably smooth as for an eruption that led the object to brighten by a factor of ~  $10^4$ . We have found that the light curve can be fitted by an exponential formula, i.e.

$$I = -2.5 \log(F_I), \qquad F_I = F_0 + F_1 \exp(t/\tau(t)), \tag{2}$$

where  $F_0 = 1.6 \times 10^{-7}$ ,  $F_1 = 2.1 \times 10^{-8}$  are fluxes in the Vega units, t = JD - 2454550, and  $\tau(t)$  is a time scale, allowed to evolve with time. The time evolution of  $\tau$ , resulting in the fit shown with the full line in the upper part of Fig. 5, is displayed in the lower part of the figure. These results show that during the initial ~5 months of the rise, i.e. up to JD ~ 2454690, the time scale was constant and equal to ~27 days. During this phase the object brightened by ~3.5 magnitudes. Then the times scale quickly decreased and the object brightened by ~6.5 magnitudes during a month.

The *I* light curve of V1309 Sco during the decline observed by OGLE-IV in 2010 is presented in Fig. 6. As can be seen from the figure, at the begining of the presented phase the decline was relatively smooth. Similar behaviour (smooth evolution) was also observed in 2009 (epoch 9 in Fig. 1). However, as the decline continues in 2010, oscillations of a few hundredths magnitude on a time scale of hours appear. A periodogram analysis shows that there is no significant periodicity in these light variations.

In a few cases during the outburst and decline multiband photometric observations are available. They allow us to estimate the spectral type and effective temperature of the object. This is the case of 7 dates in September 2008, when multiband photometry was done by the AAVSO team. On 4 and 16 October 2008 the object was observed by Rudy et al. (2008) using the Infrared Telescope Facility and *JHK* magnitudes were derived. These results can be combined with *BV* AAVSO measurements and an *I* OGLE magnitude obtained in the same time period. On 13 April 2009, as well as in a few cases in March-April 2010 and October 2010 *V* photometric measurements (together with *I*) were obtained by OGLE. On 28 August 2010 the object was observed by one of us (M.H.) with the SAAO 1.0 m. Elizabeth Telescope with the Bessell *BVRI* filters. The observations were reduced using standard procedures in the IRAF



**Fig. 5.** The *I* light curve of V1309 Sco during its rise to maximum in 2008 (upper part). Full points: data from OGLE III, open points: data from AAVSO. The line shows a fit of an exponential formula (Eq. 2). The time scale, used in the formula, is plotted in the lower part of the figure.

package. The photometry of the star was obtained with the DAOPHOT package implemented in IRAF. The resultant magnitudes are  $B = 22.21 \pm 0.10$ ,  $V = 20.42 \pm 0.13$ ,  $R = 18.84 \pm 0.10$ , and  $I = 16.67 \pm 0.10$ . Note that the latter value agrees very well with an OGLE result ( $I = 16.69 \pm 0.01$ ) obtained on the same date.

#### 6.2. Observed evolution

Taking our estimates of the distance (3.0 kpc, see Sect. 5.3.2) and reddening ( $E_{B-V} = 0.8$ , see Sect. 5.3.1) we can derive basic parameters of V1309 Sco during the outburst from multiband photometry, using the same method as in Tylenda (2005). The resultant values of the effective temperature, radius, and lu-




Fig. 6. The *I* light curve of V1309 Sco during the decline in 2010.

Table 1. Basic parametres of V1309 Sco in outburst and decline


JD	data source	sp.type	$T_{\rm eff}$	$R_{\rm eff}({ m R}_{\odot})$	L (L <sub>o</sub> )
4718.9	AAVSO	F9	5830	174.	3.16 10 <sup>4</sup>
4721.0	AAVSO	G1	5410	177.	$2.40\ 10^4$
4722.9	AAVSO	G5	4870	209.	$2.22\ 10^4$
4724.9	AAVSO	K1	4360	252.	$2.06\ 10^4$
4730.9	AAVSO	K3	4150	297.	$2.36\ 10^4$
4735.0	AAVSO	K4	3980	312.	$2.21\ 10^4$
4737.9	AAVSO	K5	3900	297.	$1.84 \ 10^4$
4750.	AAVSO/	M4	3510	155.	$3.27 \ 10^3$
	OGLE/ITF				
4934.8	OGLE	M7	3130	34.	98.5
5282.0	OGLE	M5	3370	9.5	10.4
5437.0	SAAO	M4	3420	6.9	5.9
5474.5	OGLE	M3	3565	5.4	4.2

minosity are presented in Table 1 and Fig. 7. They show that the evolution of V1309 Sco during outburst and decline was really of the same sort as those of V838 Mon (Tylenda 2005) and V4332 Sgr (Tylenda et al. 2005). In all these cases the main decline in luminosity was accompanied by a decline in the effective temperature. In later phases of the decline the objects resume a slow increase in the effective temperature. There is no doubt that V1309 Sco was not a classical nova. We know, from the present study, that its eruption resulted from a merger of a contact binary. Below we show that the energy budget of the eruption can also be well accounted for by dissipation of the orbital enregy of the progenitor.

#### 6.3. Merger powered outburst - mergeburst

The orbital energy of a binary system can be calculated from

$$E_{\rm orb} = -\frac{G M_1 M_2}{2 A} = -\frac{G (M_1 + M_2)^2}{2 A} \frac{q}{(1+q)^2}.$$
 (3)



**Fig. 7.** Evolution of V1309 during outburst and decline (see Table 1). The abscissa displays a logarithm of time in days counted from the date of the discovery.

Taking the values assumed and obtained in Sect. 5.3.2 for the V1309 Sco progenitor one gets  $E_{orb} = (0.2 - 5.6) 10^{47}$  ergs.

As derived in Sect. 5.1, the orbital period decreased by 1.2% during ~6 years. Thus the system contracted by 0.8% and its orbital energy decreased by ~  $10^{45}$  ergs during this time period,

which gives a mean rate of energy dissipation of  $\sim 10^{37}$  erg s<sup>-1</sup>. Already a small portion from this can account for the observed brightening of the object between epoch 2 and the beginning of epoch 7 (see Fig. 1). The dissipation of the orbital energy, and the accompanying loss of the orbital angular momentum, very likely resulted in mass loss from the system. This mass loss, expected to be concentrated in the equatorial plane, could have led to formation of an expanding and/or rotating disc-like envelope of the system. Since we observed the system near the equatorial plane (eclipsing binary), this envelope, when becoming optically thick, could have resulted in dimming the object for the observer. This is the likely explanation of the decline of the V1309 Sco progenitor observed during a year before the beginning of the 2008 eruption (see Fig. 1).

The progressing shortenning of the orbital period must have finally led to engulfing the secondary by the primary's envelope. This probably took place somewhere in February 2008, when the signs of the binary motion disappeared in the light curve. The secondary, or rather its core, now spiraling in the primary's, or rather common for both components, envelope started releasing the orbital energy and angular momentum at an increasing rate. The result was the gradual and relatively gentle brightenning of the object, doubling the brightness every ~19 days (Fig. 5). This phase lasted untill ~20 August (JD ~2454700), when the eruption abruptly accelarated and the object brightenned by a factor of ~300 during ~10 days. Perhaps this was the sign of the final disruption of the secondary's core deep in the primary.

As can be seen from Table 1, V1309 Sco attained a maximum luminosity of  $\sim$  3  $10^4~L_{\odot}.$  The OGLE data show that the period, when V1309 Sco was brighter than  $I \simeq 11$  magnitude, lasted 40 days. Both the maximum luminosity and the time scale of the outburst are close to theoretical expectations of a solar mass star merging with a low mass companion (Soker & Tylenda 2006). During  $\sim$ 30 days, when the luminosity of V1309 Sco was  $\gtrsim 10^4 L_{\odot}$ , the object radiated away energy of ~ 3 10<sup>44</sup> ergs. This is  $\leq 1\%$  of the energy available in the binary progenitor (see above). Of course a considerable energy was also lost during the half a year lasting rise, as well as in the decline after maximum. Certainly a significant amount was carried out in mass loss. Finally, an energy was also stored in the inflated remnant. Estimates done for V838 Mon show that the total energy involved in the outburst can be a factor of 10 - 20 greater than the energy observed in radiation (Tylenda & Soker 2006). Even than the available orbital energy can cover the budget.

Our SAAO photometry done in August 2010, as well as the OGLE V and I measurements obtained in 2010 (see Sect. 6.1), i.e about two years after the outburst maximum, show that V1309 Sco reached a luminosity (see Table 1) comparable to its preoutburst value (see Sect. 5.3.2). The object however was then significantly cooler than before the outburst (early M-type spectrum versus K1-2). The latter agrees with what was observed in V838 Mon and V4332 Sgr, displaying M-type spectra several years after their outbursts (Tylenda 2005; Tylenda et al. 2005). However the drop in luminosity of V1309 Sco was unexpectedly deep. Both V838 Mon and V4332 Sgr at present (many years after outburst) remain significantly more luminous than their progenitors. Most likely the reason for the deep decline of V1309 Sco is similar to that suggested above to explain the decline of the progenitor in 2007–8. In the merger process, especially during its initial, relatively gentle phases, say, untill August 2008, mass loss probably occured mainly in directions close to the orbital plane of the progenitor. When contracting after the outburst, the merger remnant might have largely disappeared for us behind the disc-like envelope. Moving small scale

blobs of the envelope matter, absorbing and scattering the light from the central object, could have been responsible for the short term variability seen in Fig. 6. In any case, the situation is probably similar to that observed in the V4332 Sgr remnant, where the central object is most likely hiddened in an opaque dusty disc (Kamiński et al. 2010).

#### 7. Conclusions

The principal conclusion of our study is that all the observed properties of V1309 Sco, i.e. the light curve of the progenitor during 6 years preceding the 2008 eruption, as well as the outburst itself, can be consistently explained by a merger of a contact binary. This is the first case of a direct observational evidence showing that the contact binary systems indeed end up their evolution by merging into single objects, as predicted in numerous theoretical studies of these systems.

Our study also provides a conclusive evidence in favour of the hypothesis that the V838 Mon type eruptions (red novae) result from stellar mergers, as originally proposed in Soker & Tylenda (2003) and Tylenda & Soker (2006). In particular, long and short term variabilities of the progenitors, as those of V838 Mon and V4332 Sgr (Goranskij et al. 2007; Kimeswenger 2007), which was sometimes raised as evidence against the merger hypothesis, appears now natural in view of our data for the V1309 Sco progenitor. We do not claim that all the observed eruptions of the V838 Mon type are mergers of contact binaries. There can be different ways leading to stellar mergers. What the case of V1309 Sco evidently shows is that the observational appearances of a stellar merger are indeed the same as those observed in the V838 Mon type eruptions.

The outburst of V1309 Sco was shorter and less luminous than those of V838 Mon and the extragalactic red novae. The latter objects attained luminosities of about or above  $10^6~L_{\odot}$  and their eruptions lasted a few months. These differences are most likely due to the masses of merging stars. In the case of V838 Mon an  $\sim~8~M_{\odot}$  primary was probably involved (Tylenda & Soker 2006) rather than a  $\sim~1~M_{\odot}$  one of V1309 Sco.

V1309 Sco, a rather overlooked object (only one research paper published so far, i.e. Mason et al. 2010), deserves much more attention of the observers and astrophysicists, similarly as the other V838 Mon type objects. Apart from supernovae, they belong to the most powerful stellar cataclysms. As often happens in the nature, cataclysms destroy old worlds, but also give birth to new ones. What comes out from the stellar mergers? Fast rotating giants, similar to FK Com? Peculiar stars with circumstellar discs, when new generation planets can be formed? To unswer these questions we have just to follow the evolution of the V838 Mon type objects, V1309 Sco in particular.

Acknowledgements. The OGLE project has received funding from the European Research Council under the European Community's Seventh Framework Programme (FP7/2007-2013) / ERC grant agreement no. 246678. The research raported on in this paper has partly been supported by a grant no. N N203 403939 financed by the Polish Ministery of Sciences and Higher Education. The authors are very grateful to W. Dziembowski and K. Stępień for their valuable comments and suggestions when discussing various scenarios interpreting the light curve of the V1309 Sco progenitor. The periodigram analysis described in this paper was done using a ZUZA code (version 1.2/rev5) of Alex Schwarzenberg-Czerny. This paper also uses observations made at the South African Astronomical Observatory (SAAO).

#### References

Berger, E., Soderberg, A. M., Chevalier, R. A. et al. 2009, ApJ, 699, 1850 Bond, H. E., Henden, A., Levay, Z. G. et al. 2003, Nature422, 405

- Bonnell, I. A., Matthew, R. B. & Zinnacker, H. 1998, MNRAS, 298, 93
- Bopp, B. W. & Stencel, R. E. 1981, ApJ, 247, L131
- Eggleton, P. P. 1983, ApJ, 268, 368
- Girardi, L., Bressan, A., Bertelli, G., & Chiosi, C. 2000, A&AS, 141, 371
- Goranskij, V. P., Metlova, N. V., Shugarov, S. Yu. et al. 2007, in "The Nature of V838 Mon and its Light Echo", R. L. M. Corradi & U. Munari eds., ASP Conf. Ser., 363, 214
- Iben, I., Tutukov, A. V. 1992, ApJ, 389, 369
- Jiang, D., Han, Z., Wang, J., Jiang, T. & Li, L. 2010, MNRAS, 405, 2485
- Kamiński, T, Schmidt, M., & Tylenda, R. 2010, A&A, 522, A75
- Kashi, A., Frankowski, A., & Soker, N. 2010, ApJ, 709, L11 Kimeswenger, S. 2007, in "The Nature of V838 Mon and its Light Echo", R. L. M. Corradi & U. Munari eds., ASP Conf. Ser., 363, 197
- Korhonen, H., Berdyugina, S. V., Hackman, T., Ilyin, I. V., Strassmeier, K. G. & Tuominen, I. 2007, A&A, 476, 881
- Kulkarni, S. R., Ofek, E. O., Rau, A. et al. 2007, Nature, 447, 458
- Lawlor, T. M. 2005, MNRAS, 361, 695
- Leonard, P. J. T. 1989, AJ, 98, 217
- Martini, P., Wagner, R. M., Tomaney A, et al. 1999, AJ, 118, 1034
- Mason, E., Diaz, M., Williams, R. E., Preston, G. & Bensby, T. 2010 A&A, 516, A108
- Mould, J., Cohen, J., Graham, J. R. et al. 1990, ApJ, 353, L35
- Munari, U., Henden, A., Kiyota, S. et al. 2002, A&A, 389, L51
- Munari, U., Henden, A., Vallenari, A. et al. 2005, A&A, 434, 1107
- Nakano, S. 2008, IAU Circ., 8972
- Paczyński, B., Szczygieł, D. M., Pilecki, B. & Pojmański, G. 2006, MNRAS, 368, 1311
- Rasio, F. A. 1995, ApJ, 444, L41
- Robertson, J. A. & Eggleton, P. P. 1977, MNRAS, 179, 359
- Rudy, R. J., Lynch, D. K., Russell, R. W., Sitko, M., Woodward, C. E., & Aspin, C. 2008, IAU Circ., 8997
- Schwarzenberg-Czerny, A. 1996, ApJ, 460, L107
- Shara, M. M., Yaron, O., Prialnik, D., Kovetz, A., Zurek, D. 2010, ApJ, in press, arXiv:1009.3864
- Soker, N. & Tylenda, R. 2003, ApJ, 582, L105
- Soker, N. & Tylenda, R. 2006, MNRAS, 373, 733 Soker, N. & Tylenda, R. 2007, in "The Nature of V838 Mon and its Light Echo",
- R. L. M. Corradi & U. Munari eds., ASP Conf. Ser., 363, 280
- Tylenda, R. 2005, A&A, 436, 1009
- Tylenda, R., Crause, L. A., Górny, S. K., & Schmidt, M. R. 2005, A&A, 439, 651
- Tylenda, R. & Soker, N. 2006, A&A, 451, 223
- Udalski, A. 2003, Acta Astron., 53, 291
- Udalski, A., Szymański, M. K., Soszyński, I. & Poleski, R. 2008, Acta Astron., 58.69
- Yakut, K. & Eggleton, P. P. 2005, ApJ, 629, 1055

# Appendix A: Periodograms from the preoutburst observations

Figures A.1 and A.2 present periodograms derived from the OGLE observations of the progenitor of V1309 Sco in 2002 - 2007. In all the epochs the dominating peaks is at a frequency of  $\sim 0.7 \text{ day}^{-1}$ , which we interpret as a frequency of the orbital period of the contact binary progenitor (Sect. 5.2). In earlier epochs, there is also a strong peak at a frequency of  $\sim 1.4 \text{ day}^{-1}$ , which is due to the fact that the observations can be also reasonably fitted with quasi-sinusoidal variations having a period twice shorter than the orbital period. In 2006 this peak is much weaker and practically disappears in 2007, which is due to the evolution of the light curve, as displayed in Fig. 3. All the other peaks are aliases resulting from combinations of the two above frequencies and a frequency of  $1 \text{ day}^{-1}$ .





Fig. A.1. Periodograms from the observations of the progenitor of V1309 Sco in 2002-2004.



**Fig. A.2.** Periodograms from the observations of the progenitor of V1309 Sco in 2005–2007.